

# AMOXXY 150 L.A.

## INJECTABLE SUSPENSION

**Composition:**

Each ml suspension contains: 150 mg amoxycillin base (as trihydrate)

**Spectrum of activity:**

Amoxycillin is a broad spectrum, semi-synthetic penicillin with activity against both Grampositive and Gramnegative bacteria. Sensitive are streptococci, *E. coli*, *Pasteurella* spp., *Proteus* spp., *Salmonella* spp., *Corynebacterium* spp., staphylococci ( $\beta$ -lactamase negative), *Clostridium* spp.

The drug is destroyed by e.g. penicillinase-producing staphylococci.

Some Gramnegative strains are resistant against amoxycillin (R-

plasmid). After intramuscular injection amoxycillin is quickly resorbed without irritation at the injection site.



**Indications:**

Infections of the alimentary tract, respiratory tract, urogenital tract, coli-mastitis and secondary bacterial infections during the course of viral diseases, in horses, cattle, pigs, sheep, dogs and cats.

**Contra-indications:**

New-borns, small herbivores (such as guinea pigs, rabbits), hypersensitivity to penicillins, renal dysfunctions, infections caused by penicillinase-producing bacteria.

Amoxycillin is incompatible with fast-acting bacteriostatic antimicrobial drugs (e.g. tetracyclines, chloramphenicol, sulfonamides).

**Side-effects:**

Hypersensitivity reactions may occur, e.g. anaphylactic shock.

**Dosage and administration:**

The general dose is 15 mg/kg bodyweight i.m. (this is achieved by injection of 1 ml per 10 kg b.w.), daily during 3 consecutive days.

In severe cases this dose may be given twice daily, (as there are no problems of toxicity).

Shake well before use.

**Withdrawal time:**

For meat: 7 days.

For milk: 3 days.

**Storage:**

Store between 8 °C and 15 °C and protected from light.

**Shelf life:**

2 years.

**Presentation:**

100 ml multidose vial.

### FOR VETERINARY USE ONLY