

FLOXATRIL 10 %

INJECTABLE SOLUTION

Composition:

Each ml solution contains: 100 mg norfloxacin.

Spectrum of activity:

Norfloxacin, a quinolone derivative from the third generation, has antimicrobial activity against a wide variety of aerobic Gramnegative and Grampositive bacteria, including *Pseudomonas aeruginosa*, *Neisseria gonorrhoeae*, members of the family *Enterobacteriaceae*, *Staphylococcus aureus*, *Actinobacter* spp., *Haemophilus* spp., *Salmonella* spp., *Bordetella bronchiseptica* and *Pasteurella multocida*. Norfloxacin is also active against veterinary mycoplasmas.



Indications:

Norfloxacin is highly effective in the treatment of uncomplicated and complicated urinary tract infections, gonococcal urethritis, mastitis, salmonellosis, respiratory infections including enzootic pneumonia in cattle, pigs, dogs and poultry.

It is also used as part of selective digestive tract decontamination in intestinal diseases.

Contra-indications:

Avoid using with other medications.

Side effects:

High doses can cause nephrotoxicity and crystalluria.

Dosage and administration:

Administration by subcutaneous or intramuscular injection. For large animals (horse, cattle, sheep) and poultry both therapeutic and prophylactic daily dose is 2.5 – 5.0 mg/kg bodyweight (1 ml per 20-40 kg bodyweight) for 3 – 5 days.

Withdrawal time:

For meat: 4 days.

For eggs: 4 days.

For milk: 4 days.

Storage:

Store at room temperature and protected from light.

Shelf life:

2 years.

Presentation:

30 ml and 100 ml multidose vial.

FOR VETERINARY USE ONLY