

ISOPHEN

INJECTABLE SOLUTION

Composition:

Each ml solution contains: 120 mg phenylbutazone sodium, 240 mg isopropylaminophenazone.

Activity:

Phenylbutazone is a synthetic non-steroidal anti-inflammatory agent which also exhibits analgesic and antipyretic properties (without loss of consciousness). It acts symptomatically. Phenylbutazone is widely used to treat or to give symptomatic relief from painful arthritis and skeletal muscular disorders. Furthermore it is frequently beneficial in treatment of soft tissue or nonarticular rheumatism.

Isopropylaminophenazone also belongs to the non-steroidal anti-inflammatory drugs, and has the same properties as phenylbutazone, but at the same time it decreases the normal toxicity of phenylbutazone, as a lower dose of phenylbutazone will be sufficient.



Indications:

All conditions which are accompanied with pain or inflammation and particularly when these conditions are accompanied with oedema, e.g. (poly)arthritis, peri-arthritis, tendinitis, tendovaginitis, bursitis, neuritis, myositis, enchondrosis intervertebralis with or without paresis, meningitis, distorsions, colics, (enzootic) pneumonia and (in combination with antibiotics) mastitis.

Contra-indications:

Administration to animals with severe cardiac-, renal- or hepatic injury, newborn animals, combination with strong plasma-protein binding medicines (e.g. salicylates or sulfonamides).

Caution:

Do not administer intramuscularly to horses.

Stop treatment in racing-horses at least 5 days before the race (dope).

Intramuscular administration of large doses results in tissue damages, therefore avoid administration of more than 2 ml (dogs) or 10 ml (cattle) per injection site.

Dosage and administration:

For intramuscular, subcutaneous or slowly intravenous administration.

Horses: 10-20 ml intravenously per animal.

Cattle: 10-20 ml i.m. or i.v. per animal.

Calves, foals: 4-6 ml i.m. or s.c. per animal.

Pigs: 5-12 ml i.m. or s.c. per animal.

Piglets : 0.5-1.5 ml i.m. or s.c per animal.

Dogs: 0.5-5 ml i.m. or i.v. per animal.

Cats: 0.2-0.5 ml i.m. or i.v. per animal.

Withdrawal time:

For milk: 4 days. For meat: 12 days

Storage:

Store between 8 C and 15 C and protected from light.

Presentation:

100 ml multidose vial.

FOR VETERINARY USE ONLY