

LIDOCAINE 20

INJECTABLE SOLUTION

Composition:

Each ml contains:
20 mg lidocaine hydrochloride
0.025 mg noradrenaline bitartrate

Activity:

Lidocaine is a local anesthetic.
Noradrenaline causes vasoconstriction, resulting in a prolonged anesthetic effect of the lidocaine and a decreased chance of side-effects.



Indications:

Local anesthetic for local-, infiltration-, conduction- and lumbal anesthesia.

Contra - indications:

Intravenous administration.

Side - effects:

When used at high doses somnolence may occur.

Dosage and administration:

Nerve block or conduction block:

diagnostic injections in horse and cow: 5-10 ml.
anesthesia of the cornual nerve of the cow (dehorning): 8-10 ml each side.

Infiltration-anesthesia:

e.g. sectio caesaria (standing animal): maximal 100 ml.

Sacral anesthesia:

for surgery in standing animals:

horse: 5-10 ml (slow administration)
cow: 4-6 ml (slow administration)
sheep: 2-3 ml (slow administration).

Epidural anesthesia:

for surgery in the lying animal:

horse: 25-75 ml (slow administration)
cow: 15-60 ml (slow administration)
small animals: 1.5 ml per 10 cm length of the vertebral column (slow administration).

Paravertebral block (cow):

10 ml each nerve root.

Withdrawal time:

For meat: 1 day.
For milk: 1 day.

Storage: Store between 8 °C and 15 °C and protected from light.

Presentation: vial of 100 ml.

FOR VETERINARY USE ONLY