

PANTRISUL

INJECTABLE SOLUTION

Composition:

Each ml solution contains: 40 mg trimethoprim.
200 mg sulfamethoxazole.

Spectrum of activity:

The combination of trimethoprim and sulfamethoxazole acts synergistic and in many cases bactericidal against a large number of Gram-positive and Gram-negative bacteria, e.g. some staphylococci, streptococci, *E. coli*, *Actinomyces* spp., *Actinobacillus pleuropneumonia*, *Klebsiella* spp., *Listeria monocytogenes*, *Salmonella* spp., *Proteus mirabilis*, *Pasteurella* spp., *Enterobacter aerogenes*, *Haemophilus influenza* and *Corynebacterium* spp..

**Indications:**

Infections caused by micro-organisms susceptible to trimethoprim and/or sulfamethoxazole (like respiratory-, alimentary- and urogenital tract infections, secondary bacterial infections (after viral infections), foot rot, mastitis, arthritis and phlegmones), in horses, cattle, pigs, dogs and cats.

Contra-indications:

Serious liver- or kidney insufficiencies, aciduria, hypersensitivity to sulphonamides.
The combination is incompatible with para-aminobenzoic acid and esters of it (procaine, tetracaine). Don't administer simultaneously with hexamethylene tetramine (methenamine).

Side-effects:

Occasionally, temporary swellings at injection sites may occur (for this reason subcutaneous injections should be avoided).
Folic acid deficiency, crystalluria and/or blood dyscrasias may occur, especially if treatment is prolonged.
Hypersensitivity and cross-hypersensitivity reactions to sulphonamides.

Dosage and administration:

For intramuscular or slow intravenous (horses, foals) injection.
The general dose is: 1 ml per 10-15 kg bodyweight, once daily, during 5 days (if necessary).

Withdrawal time:

For meat: 8 days.
For milk: 4 days.

Storage:

Store between 15 °C and 25 °C and protected from light.

Shelf life:

2 years.

Presentation:

100 ml multidose vial.

FOR VETERINARY USE ONLY