


# S U L F A D I M I N E 3 3 3

## INJECTABLE / ORAL SOLUTION

### Composition:

Each ml solution contains: 333 mg sulfadimidine sodium.



### Spectrum of activity:

Sulfadimidine is a sulphonamide with antimicrobial activity against Gram-positive cocci (hemolytic streptococci, pneumococci and some staphylococci), against some Gram-negative bacteria (like *Pasteurella* spp., *E. coli*, *Salmonella* spp.) and against some *Rickettsiae* and protozoa. Sulfadimidine is rapidly absorbed from the site of application and slowly excreted by the organism, so in most animals a single daily dose is sufficient. Bacterial resistance against sulphonamides mostly is permanent and cross-resistance between sulphonamides is common.

### Indications:

Pneumonia, foot rot, diphtheria, actinobacillosis, colibacillosis, salmonellosis, pasteurellosis, enteritis, polyarthritis, streptococcal mastitis, dysentery, lymphangitis, young's septicemia and coccidiosis in horses, cattle, pigs, sheep, goats, dogs and poultry.

### Contra-indications:

Liver or renal dysfunctions, diseases with decreased urine production (oliguria, anuria), hypersensitivity to sulphonamides. Sulfadimidine is incompatible with para-aminobenzoic acid and its esters (e.g. procaine or tetracaine). Don't administer simultaneously with hexamethylene tetramine (methenamine).

### Side-effects:

Sometimes hemorrhagic enteritis (in cattle), kidney damage, hypersensitivity reactions or crystalluria.

### Dosage and administration:

For slow intravenous, intramuscular or oral administration.  
Parenteral administration. Cattle, sheep, goats, pigs, horses: 3 ml per 10 kg b.w., during 3 – 5 days.  
Oral administration. Poultry: 5 – 6 ml per litre of drinking water, during 3 – 6 days. Cattle: 3 ml per 10 kg b.w., during 3 – 6 days. Provide ample drinking water during treatment.

### Withdrawal time:

For milk: 5 days.  
For meat: 8 days (horse, cow, sheep, goat)  
10 days (calf, pig).

### Storage:

Store at room temperature and protected from light. Keep out of reach of children

### Shelf life:

2/3 years.

### Presentation:

100 multidose vial. Plastic vial 250 ml and 500 ml

## FOR VETERINARY USE ONLY