

TYLOCARE (TYLOSIN 200)

INJECTABLE SOLUTION

Composition:

Each ml solution contains:
200 mg tylosin base (as tartrate).

Spectrum of activity:

Tylosin, a macrolide antibiotic, is active against particularly Gram-positive bacteria like *Streptococcus* spp. and *Staphylococcus* spp. (including *Staphylococcus aureus*), some Spirochetes (including *Leptospira*); *Actinomyces*, Mycoplasmas (PLO), *Haemophilus pertussis*, *Moraxella bovis* and some Gram-negative cocci.

After parenteral administration, therapeutically active blood concentrations of tylosin are reached within 2 hours.

Indications:

Infections, caused by micro-organisms susceptible to tylosin, like e.g. respiratory tract infections in cattle, sheep and pigs and poultry, Dysentery Doyle in pigs, dysentery and arthritis caused by Mycoplasmas, mastitis and endometritis.

Contra-indications:

Hypersensitivity to tylosin, cross-hypersensitivity to macrolides. Use in laying hens.

Side-effects:

Sometimes, local irritation at the injection site may occur (inject slowly).

Dosage and administration:

For intramuscular or subcutaneous administration.

The general dose is: 1 ml per 20 kg bodyweight daily, during 3-5 days.

Withdrawal time:

For meat: 8 days (cattle).

4 days (pigs, poultry).

For milk: 4 days.

Storage:

Store between 15 °C and 25 °C and protected from light.

Shelf life:

2 years.

Presentation:

100 ml and 250 ml multidose vial.



FOR VETERINARY USE ONLY