

FLUNIMEG 50

INJECTABLE SOLUTION

Composition:

Each ml solution contains: 50 mg flunixin (as meglumine).

Spectrum of activity:

Flunixin meglumine is a non-steroidal drug, possessing analgesic, anti-inflammatory and antipyretic properties. Active blood concentrations are reached within 15-60 minutes after injection and last for 30-40 hours.

Indications:

Cattle : inflammation associated to diseases of the respiratory tract.
Horses : visceral pains associated to colic. Inflammations and pains associated to musculoskeletal disorders.

Contra-indications:

Renal dysfunctions.

Side-effects:

Gastro-intestinal disturbances like gastric ulcer and occult bleeding.
Nephrotoxicity or changes in renal function (papillary necrosis).

Dosage and administration: For intravenous administration:

Cattle: 2.2 ml FLUNIMEG 50 per 50 kg bodyweight (2.2 mg flunixin per kg bodyweight), daily, for maximum 3 consecutive days

Horses: 1.1 ml FLUNIMEG 50 per 50 kg bodyweight (1.1 mg flunixin per kg bodyweight), daily, for maximum 5 consecutive days

Opened vials should be used within 28 days.

Withdrawal time:

Cattle : Meat: 7 days. Milk: 1 days

Horses : Meat 28 days.

Storage:

Store between 15 °C and 25 °C and protected from light.

Shelf life:

2 years.

Presentation:

100 ml multidose vial.



FOR VETERINARY USE ONLY